REFERENCE

Alvi, A. S. 1994. Adventitious toxins in plant origin feedstuffs:qualification and tolerance level in fish. Master's dissertation. Aligarh Msulim University, Aligarh, India.

AOAC(Association of Official Analytical Chemists). 1995. Official methods of analysis of the Assocition Official Analytical Chemists,16th edition.AOAC, Inc., Arlington,Virginia,USA

APHA. 1998. Standard methods for the examination of water and wastewater,20th edition.*in* L.S.Clesceri,A.E.Greenberg,and A.D.Eaton,editors.American Public Health Assocition, American Water Works Association, Water eEnvironment Federation,Washington,DC,USA.

Baeverfjord, G., T. Asgard, and K. D. Shearer. 1998. Development and detection of phosphorus deficiency in Atlantic salmon,*Salmo salar* parr and post-smolts. Aquaculture Nutrition 4:1-11.

Baruah, K., A. K. Pal, N. P. Sahu, K. K. Jain, S. C. Mukherjee, and D. Debnath. 2005. Dietary protein level,microbial phytase,citric acid and their interactions on bone mineralization of *Labeo rohita* (Hamilton) juveniles, Aquaculture Research 36:803-812

Cheng, Z. J. and R. W. Hardy. 2002. Effect of microbial phytase on aparent nutrient digestibility of barley,canola meal,wheat and wheat middlings, measured in vivo using rainbow trot, *Oncorhynchus mykiss.* Aquaculture Nutrition 8:271-277.

De Silva, S. S. and T. A. Andersons. 1995. Fish nutrition in aquaculture. Chapman and Hall aquaculture series 1.Chapman and Hall, London,UK.

Debnath, D., A. K. Pal, N. P. Sahu, K. K. Jain, S. Yengkokpam, and S.C.Mukherjee. 2005a. Effect of dietry microbial phytase supplementation on growth and nutrient digestibility of *Pangasius pangasius* fingerlings.Aquaculture Research 36:180-187.

Debnath, D., N. P. Sahu, A. K. Pal, K. K. Jain, S. Yengkokpam, and S.C.Mukherjee. 2005b. Mineral status of *Pangasius pangasius*(Hamilton) fingerlings in relation to supplemental phytase: absorption, whole body and bone mineral content. Aquaculture Research 36:326-335.

Duncan, D. B. 1955. Multiple range and multiple F tests. Biometrics 11:1-42.

Forster, I., D. A. Higgs, B. S. Dosanjh, M. Rowshandeli, and J. Parr. 1999. Potential for dietary phytase to improve the nutritive value of canola protein concentrate and decrease phosphorus output in rainbow trout, *Oncorhynchus mykiss* held in 11 C fresh water. Aquaculture 179:109-125.

Francis, G., H. P. S. Makkar, and K. Becker. 2001. Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish. Aquaculture Research 199:197-227.

Furuya, W. M., G. S. Goncalves, V. R. B. Furuya, and C. Hayashi. 2001. Phytase as feeding for Nile tilapia,*Oreochromis niloticus.* Perfomance and digestibility.Revista Brasileira de Zootecnia 30:924-929.

Gatlin, D. M., III and H. F. Phillips. 1989. Dietary calcium,phytate and zinc interactions in channel catfish.Aquaculture 79:259-266,

Grant, G. 1989. Antinutritional effects of soybean:a review.Progress in Food Nutritional Science 13:317-348.

Hardy, R. W. and F. T. Barroes. 2002. Diet formulation and manufacture.Pages 505-600 *in* J.E. Halver and R.W.Hardy,editors.Fish nutrition,3rd edition.Academic Press,San Diego,California,USA.

Hendricks, J. O. and G. S. Baliey. 1989. Adventitious toxins. Pages 606-644 *in* J.E. Halver, editor. Fish nutrition.Academic Press, Inc.,San Diego,Calfornia,USA.

Hossain, M. A. and K. Januccy. 1991. The effect of varying dietary phytic acid,calcium and magnesium levels on the nutrition of common carp,Cyprinus carpio. Fish nutrition in practice.Proceedings of the 4th International Symposium on Fish Nutrition and feeding, Biarritz,France.INRA,Paris,France.

ICLARM (International center for Living Aquatic Resources Management). 2001. Gentic improvement of crap spcies in Asia:final report.Asian Development Bank Regional Technical Assistance No.5711.International Center for Living Aquatic Resources Management,Penang,Malaysia.

Jackson, L. S., M. H. Li, and E. H. Robinson. 1996. Use of microbial phytase in channel catfish,*Ictalurus punctatus* diets to improve utilization of phytate phosphorus.Journal of the World Aquaculture Society 27:309-313.

Jhingran, V. G. 1991. Fish and fisheries of India. Hindustan Publication Company, New Delhi,India.

Knuckles, B. E., D. D. Kuzmicky, M. R. Gumbmann, and A. A. Betschart. 1989. Effect of myo-inositol phosphate esters on *in vitro* and in vivo digestion of protein.Journal of Food Science 54:1348-1350.

Liebert, F. and L. Portz. 2005. Nutrient utilization of Nile tilapia,*Oreochromis niloticus* fed plant based low phosphorus diets suoolemented with graded levels of different sources of microbial phytase.Aquaculture 248:111-119.

Masumoto, T., B. Tamura, and S. Shimeno. 2001. Effects of phytase on bioavailability of phosphorus in soybean meal-based diets for Japanese flounder, *Paralichthys olivaceus* .Fisheries Science 67:1075-1080.

Miranda, E. C., A. C. Pezzato, L. E. Pezzato, C. F. Graner, G. J. Rosa and L. G. Q. Pinto. 2000. Availability of calcium:phosphorus ratio in diets for Nile tilapia, *Oreochromis niloticus.*Revista Brasileira De Zootecnia 2962-2171.

O'Dell, B. L. and A. De Boland. 1976. Complexation of phytate with proteins and cation in corn germ and oliseeds meals.Journal of Agricultural and Food Chemistry 24:804-808.

Ogino, C., L. Takeuchi, H. Takeda, and T. Watanbe. 1979. Availability of dietary phosphorous in carp and rainbow trout. Bulletin of the Japanese Society of Scientific Fisheries 45:1527-1532.

Oliva-Teles, A., J. P. Pereira, A. Gouvein, and E. Gomes. 1998. Utilization of diets supplemented with microbial phytase by seabass,*dicentrarchus labrax* juvenlies.Aquatic Living Resource 11:255-259.

Papatryphon, E. and J. H Soares. 2001. The effect of phytase on apparent digestibility of four parctical plant feedstuffs fed to striped bass,*Morone saxatilis.* Auaculture Nutrition 7:161-167.

Papatryphon, E., R. A. Howell, and J. H. Soares. 1999. Growth and mineral absorption by striped bassd,*Morone saxatilis* fed a plant feedstuff bassd dite supplemented with phytase.Journal of world Aquaculture Socity 30:161-173.

Pointillart, A., A. Fourdin, and N. Fontaine. 1987. Importance of cereal phytase activity for phytate phosphirus utilization by growing piogs fed diets containing triticale or corn.Journal of Nutrition 29:907-912.

Portz, L. and F. Liebert. 2004. Growth,nutrient utilization and parameters of mineral metabolism in Nile tilapia,*Orechromis niloticus* fed plant based dites with graded levels of microbial phytase.Journal of Animal physiology and Animal nutrition 88:311-320.

Richardson, N. L., D. A. Higgs, R. M. Beames, and J. R. McBride. 1985. Influence of dietary calcium,phosphorus,zinc and sodium phytate level on cataract incidence,growth, and histolopathology in juvenile chinook salmon,*Oncorhynchus tshawytscha.*Journal of Nutrition 115:553-567.

Robinson, E. H., M. H. Li, and B. B. Manning. 2002. Comparison of microbial phytase and dicalcium phosphate for growth and bone mineralization of pond-raised channel catfish,*Ictalurus punctatus.*Journal of Applied Aquaculture 12:81-88.

Sajjadi, M. and S. G. Carter. 2004. Dietary phytase supplementation and the utilisation of phosphorus by Atlantic salmon, *salmon salar* fed a canola-meal-based dite. Aquaculture 240:417-431.

Satoh, S., W. E. Poe, and R. P. Wilson. 1989. Effect of supplemental phytate and/or tricalcium phosphate on weight gain,feed efficiency and zine content in vertebrae of channel catfish *Ictalurus punctatus.* Aquaculture80:155-161.

Schafer, A., W. M. Koppe, K. H. Meyer-Burgdorff, and K. D. Gunther. 1995. Effect of a microbial phytase on the utilization of native phosphorus by crap in a dite based on soybean meal. Water Science and Technology 31:149-155.

Simons, P. C. M., H. A. J. Versteegh, A. W. Jongbloed, P. A. Kemme, P. Slump, K. D. Bos, W. G. E. Wolters, R. F. Beudeker, and G. J. Verschoor. 1990. Improvment of phosphorus availability by microbial phytase in broilers and pigs.British Journal of Nutrition 64:525-540.

Singh, M. and A. D. Krikorian. 1982. Inhibition of trypsin activity *in vitro* by phytase.Journal of Agricultural and Food Chemistry 30:799-800.

Storebakken, T., K. D. Shearer, and A. J. Roem. 1998. Availability of protein, phosphorus and other elements in fish meal, soy-protein concentrate and phytasetreated soy-protein –concentrate-based diets to atlantic salmon,*salmo salar.*Aquaculture 161:365-379.

Sugiura, S. H., J. Gabaudan, F. M. Dong, and R. W. Hardy. 2001. Dietary microbial phytase supplementation and the utilization of phosphorus,trace minerals and protein by rainbow trout,*Oncorhynchus mykiss* fed soybean meal-based diets. Aquaculture Research 32:583-592.

Teskeredzic, Z., D. A. Higgs, B. S. Dosanjh, J. R. McBride, R. W. Hardy, R. M. Beanes, M. Simell, T. Vaara, and R. B. Bridges. 1995. Assessment of unphytinized and dephytinized rapseed protein concentrate as sources of dietary protein for juvenlie rainbow trout,*oncorhynchus mykiss.* Aquaculture131:261-277.

Usmani, N. and A. K. Jafri. 2002. Influence of dietary phytic acid on the growth, conversion efficiancy,and carcass composition of mrigal,*Cirrhinus mrigala* fry. Journal of World Aquaculture society 33:199-204.

Van den Ingh, T. S. G. A. M., A. Krogdahl, J. J. Oil, H. G. C. J. M. Hendricks, and J. G. J. F. Koninkx. 1991. Effect of soybean containing diets on the proximal and distal intestine in Atlantic salmon,*Salmo salar. A morphological study.* Aquaculture 94:297-305.

Van Weerd, J. H., K. H. A. Khalaf, F. J. Aartsen, and P. A. T. Tijssen. 1999. Balance trials with african catfish,*Clarias gariepinus* fed phytase-treated soybean meal-based diets. Aquaculture Nutrition 5:135-142.

Vielma, J. and S. P. Lall. 1998. Phosphorus utilization by Atlantic salmon,*salmo salar* reared in freshwater is not influenced by higher dietary calcium intake. Aquaculture60:117-128.

Vielma, J., S. P. Lall, J. Koskela, F. J. Schoner, and P. Mattilla. 1998. Effects of dietary phytase and cholecalciferol on phosphorus bioavailability in rainbow trout,*Oncorhynchus mykiss.*Aquaculture 163:309-323.

Vielma, J., T. Makinen, P. Ekholm and J. Koskela. 2000. Influence of dietary soy and phytase levels on performance and body composition of large rainbow trout,*Oncorhynchus mykiss* and algal availability of phosphorus load.Aquaculture 183:349-362.

Vielma, J., K. Ruohonen, and M. Peisker. 2002. Dephytinization of two soy proteins increases phosphorus and protein utilization by rainbow trout,*Onchorhynchus mykiss.*Aquaculture 204:145-156.

Vielma, J., K. Rouhonen, J. Gabaudan, and K. Vogel. 2004. Top spraying soybean meal-based diets with phytase improves protein and mineral digestibilities but not Lysine utilization in ranibow trout,*Oncorhynchus mykiss.*Aquaculture Research 35:955-964.

Yan, W. and R. C. Reigh. 2002. Effects of fungal phytase on utilization of dietary protein and minerals,and dephosphorylation of phytic acid in the alimentary tract of channel catfish,*Ictalurus punctatus* fed an all-plant- protein diet.Journal of World Aquaculture Society 33:10-22.

Yi, Z. and E. T. Kornegay. 1996. Sites of phytase activity in gastrointestinal tract of young pigs.Animal Feed Science Technology 61:361-368.

Yoo, G. Y., X. Wang, S. Choi, K. Han, J. C. Kang, and S. C. Bai. 2005. Dietary microbial phytase increased the phosphorus digestibility in juvenile Korean rockfish,*Sebastes schlegeli* fed diets containing soybean meal.Aquaculture 243:315-322.

Zongjia, J., Z. J. Cheng, and R. W. Hardy. 2003. Effects of extrusion and expelling processing, and microbial phytase supplementation on apparent digestibility coefficients of nutrients in full-fat soybeans for rainbow trout,*Oncorhynchus mykiss.*Aquaculture 218:501-514. APHA, 1989. Stadard methods for the exzmination of water and waste water 17th Edition (American Public health Association) Washington.

AOAC (1990). Official methods of analysis (15th ed.) – K.Heldrich (ed) . Association of Official Analytical Chemists. Virginia. U.S.A p.1298 Banerjee,S.C and Bagchi, M.M.(1970). A note in seasonal changes of fat in Rohu, *Labeo rohita* (Ham). *J. inland Fish. Soc. Ind*. **1**: 139-142.

Bhimachar, B.S (1971). Food of the carp hatching in fish culture. *Indian Fmg.***20** (12):45-46.

Brooks, J.K. and Dodson, S.I;1965. Science, 150:28 - 35.

Brown, M.E. 1974. Seed and root bacterization Annce. Rev. phytopathol. **12** :181- 197

Boyd, C.E. Hollerman, W.D.plumb.J.A; Saeed,M.; 1984. Effect of treatment with a commercial bacterial suspension on water quality in channel catfish ponds. Prog. Fish cult. 46, 36-40.

Blackburn,T.H. and K. Henriksen.1985. nitrogen cycling in different types of sediments from Danish water, Limnol.Occanogr.**28** (3);477-493.

Cooper, R. 1959. Bacterial fertilizers in the Soviet Union. Soils Fert. **22**: 327-333.

Das, S. (1949). The relative importance oforganic manures and inorganic fertilizers in tropical soils. *Indian J.agric. sci.,***19** (1) .,41-49.

Dayal, P. (1982). Importance of fish culture in developing countries. *Bull. Agric. Congo Belgi.:* **17**.

Fang Xiuzheng; Guo Xianzhen and Wang Jikum. 1989. The preliminary studies on the heterotrphc bacteria in high-yeild fish ponds. Fishery Journal **13** (20 : 101-109.

Guo Xianzhen; Fang Yingxue and Wang Jikun. 1988. The preliminary studies on the bacteria types in the fish ponds applied with four kinds of animal measure and the effects upon ecosystem, and yield. J.Chinese Acad. Fish. Sci. **1**(1); 18-28

Garg, A.C.,Indani, M.A. and Abraham, T.P. (1971). Organic manures *ICAR Tech Bull (Agri)* **32**: 83p.

Ghosh, S.R. (1975). A study on the relative efficiency organic manure and the effect of salinity and its mineralization in brakish water fish farm soil. *Aquaculture*, **5** (4): 359 – 366.

Gopal Dutta, N.H., Appraisal of North America fish culture fertilization studies. FAO Fish Rep. **3** (44): 13 – 26.

Gopal Krishnan, V. and Srinath, E.G(1963). Some experimental observations on the use of activated sludge as fertilizer for fish culture. *Proc. Indian Acad Sci.***47** b(6): 379 – 388.

Gooch,B.C. (1967). Appraisal of North American fish culture fertilization studies. *FAO. Fish Rep.* **3** (44): 13 – 26.

Govind, B.V., Rajgopal, K.V, and Singh, G.S. (1978). Studies on the comparative efficiency of organic manures as fish producers. *J. Inland Fish Soc.* India **10**: 101 – 106.

Gupta, A. (1983). Exploitation and management of pisciculture respirces (a case study of Birbhum). Ph.D. thesis (unpublished) *Visva Bharati*.

Hickling, C.F. (1971) *Fish culture*. 2nd ed. London, Faber and Faber, 317.

Hora, S.L. (1946) The village pond and fish production. *Indian Fmg.* (Special number): 190 – 195.

Hora, S.L. (1951). Fish Culture in Rice field. Sci. **20** (7): 171 – 173.

Hora, S.L. and pillay, T.V.R. (1962). Hand book of fish culture in the Indo-pacific fisheries region, *FAO fish, Biol.* Tech. Pap. (14): 203.

Homell, J. and Nayudu, M.R. (1923). A contribution to the life history of the Indian sardine, with notes on the plankton of the Malabar coast. *Madras, Fish Bull,* **17**: 129-197.

Hrback, J; 1962. Rozpravy Ceskoslovenske Academic ved; **72**: 1-116

Hopkins, S.J; Hamilton, R.D.11, Sandifer, P.A; Browdy, C.L; Stokes, A.D; 1993. Effect of water exchange rate on production, water quality, effluent characteristics and nitrogen budgets of intensive shrimp pods J. World Aquacult. Soc. 24. 304-320.

Jafry, D.E. 1998. use of probiotics in penaeid shrimp growout. Aquacult. Mag. 24 (1).

62-67.

Jafry, A.K. Khawaja, D.K. and Quasim, S.Z/ (1964). *Fish Technol.* **2**: 148 – 157.

Jhingran, V.G. (1982). *Fish and Fisheries of India.* Hindustan publishing Corporation (India). Delhi, 666.

Lakshmanan, M.A.V., Sen, P.R., Murty, D.S. and Chakraborty, D.P. (1968) Preliminary study on the rearing of carp fingerlings. *Indian J. Fish.* **15**: 40 – 52.

Lovern, J.A. (1983a). Fat metabolism in fishes. XII. Seasonal changes in the composition of herring fat. *Biochem.J.***32**: 676 – 680.

Lovern, J.A. (1983b). Fat metabolism in fishes. XIII. Factor influencing the composition of the depot fat of fishes. *Biochem. J.* **32**: 1214 – 1224.

Lowry, O.H; Rosbrough, N.J; Forr, A,L. and Ramda; R.J., 1951. J. B 101, Chem. **193**: 265.

Lukowicz, M.;1982. Aquaculture Engineering,, **1**:121 – 137.

Liu Guocai: Bao wenzhong and Liu Zhenqi.1992 (a). The growth and the seasonal changes of bacteria biomass in fishpons. Fishery Journal **16** (1): 24 – 31.

Liu Guocai: Bao wenzhong and Liu Zhenqi.1992 (b). preliminary studies on the bacteria in pond mud. *J.Fish. Biology.* **16** (3): 284 - 286.

Maeda, H. 1987.H. 1987. Changes in the qualities of water and bottom sediment with the development of anoxie layer in a stratified lake. Nippon Sojima **53** (7): 1281-1288.

Mahboob, S., Sheri, A.N. (1997). Growth performance of major, common and some Chinese carps under composite culture system with special reference to pond fertilization. J. Aquacult. Trop. **12** (3): 201 – 207.

Meehean, O.L. (!933). the role of fertilizers in pond fish production and some ecological aspects. *Trans. Amer. Fish. Soc* **63**: 103 – 109.

Miah, M.S./ Alam, M.S., Neaguddin, M., Gupta, M.V., Shah, M.S. (1996). Studies on production craps under polyculture system in fermer's pond's. *Indian J. Fish.* **43** (4): 375 – 379.

Mitra, B. and Gupta, A. (1981). Effect of poultry manure growth of three Indian major craps, *All Indian Seminar on Fish Biology, Bihar University,* Muzaffarpur: Abs. 4.

Mitra, B. and Rej. S.K. (1976). Variation of musle lipid of Magur fish (*Clarias batrachus Lin.*) with respect to size, weight and reasons. Applied Nutrition, **4** : 103 – 106.

Mitra, B., Choudhury, A. and Mukherjee, K. (1977). Variation of muscle lipid in fishes. *Proc. Zool. Soc. Calcutta.* **30**:115 – 119.

Mitra, B., Gupta, A. and Laha, U. (1987). Effects of some manures on the growth and production of major craps in village ponds of district Birbhum, West Bengal *environ. Ecol.* **5**: 381 – 385.

Mortimer, C.H. and Hickling, F.C. (1954). *Fertilizers in fish ponds. Fish publ. Lond.* (5).

Mei Zhiping: Shi Zhenfeng and Zhenfeng and Zhu Xuebao. 1995. Yhe studies on the nutritional content and the release of the fish pond mud. P.29-36. The studies of the Ecosystem in the Chinese Fish Pond. Shanghai Science and Technology Press.(Editor: Zhu Xuebao and Shi Zhenfeng).

Nandeesha, M.C. (1982). Effect of three different organic manures on the growth of cultivable carps M.F.Sc. Thesis *Universityu of Agricultural Science,* Bangalore.

Nataranjan, M. and Varghese, T.J. (1978). Digested sewage sludge cakes as manure for fish culture. *Inland Fmg.***28**: 25 – 27.

Queiroz, J.F; Boyd, C.E., 1998. Effect of a bacterial inoculums in channel catfish ponds. World Aquacult. Soc. **29**: 67 – 73.

Pandey, B.N., Dallamunshi, J.S., Choubey, B.J. and Pandey, P.K. (1976). Seasonal variation in body composition in relation to breeding cycle of an air-breathing fish, *Heteropneastes fossilis* (Bloch). J. *Indian Fish Soc. Indian.* **8** : 91 – 96.

Philips, A.M. Jr. (1969). Nutrition digestion and energy utilization. In *Fish physiology,* (W.B. Hoar and D.J. Randalleds. Academic Press, New York. **1**: 391 – 423.

Prowse, G.A. (1967). A review of the methods of fertilizing warm watewr fish ponds in Asia and far best. *FAO Fish Rep* **3** (44): 7 – 12.

Peowse, G.A. (1968). The latin square fertilizer trails. Trop. *Fish cult. Res. Inst. Wkg. Pab.* (1): 28.

Rath, R.K. (1978). Role of fishery science in human health and welfare, Proc. 65th *Ind. Sci. Cong*., Advance Notes on symposia and Discussions: *Soc. Zoology, Entomology and Fishery, Focal theme.*

Ray, P. and David, A. (1969). Poultry manure as potential plankton producers in fish nurseries. *Labdev. J. Sci. Tech.* **7B**: 229 – 231.

Roy. S. K. and Mitra, B. (1978). The trend of variation of muscle lipid in two body regions of *Anabas testudineus* (Bloch). *Indian Biologist.* **10**: 45 – 48.

Rimon, Aand Shilo, M; 1982. Bamidgeh, 34: 87 – 100.

Saha, G. N. Chattarjee, D. K. and Raman, K. (1974). Observation on manuring nursery ponds with successful rearing of carps spawn. *J. Inland Fish Soc. India* **6**: 162 – 166.

Saha, G.N. , Chatterjee, D.K., Selvaraj, C. and Mazumdar, N.M. (1978). A record of increased fish production in freshwater ponds by use of fertilizers alone. *Sci. Cult.,* **44** (9): 422 – 424.

Saha,G.N., Raman, K., Chatterjee, D.K., and Ghosh, S.R.(1975). Relative response of three nitrogenous fertilizers in different pond soil in relation to primary productivity,. Plankton and survival and growth of *Labeo rohita spawn. J. Inland Fish Soc. India.***7**: 162 – 171.

Schaperclaus, W. Die Karpfenteichswirt Schaft. (1959). *In Der Deutsche Demo Kratischen Pepublik. Sita Ber. Bd. VII. Hrt. 7 Deutsche Akad. Landwirtschaftwiss. Berlin.*

Sinha, V.R.P., Gupta, M.V., Banerjee, M.K. and Kumar, D.(1973). Composite fish culture at Kalyani, West Bengal. J. *Inland Fish Soc. India*. **5**: 201 – 207.

Smith, E.V. and Swingle, H.S. (1939). The relationship between plankton production and fish producti9on in ponds. *Trans.Amer. Fish. Soc.***68**: 309 – 315.

Sreenivasan, A.and Soundara Raj, R.(1962). Variation in the chemical composition of a freshwater fish, *Labeo Fimbriatus, Madras J.Fish. I*: 126 – 128.

Sukumaran, K.K.(1976). Progress Repport on the Karnal sub-center of Haryana for the period of November, 1972 to August, 1975. *In Third workshop on the All India Co-ordinated Research Project on Composite Fish Culture and Fish seed production.FIFRI.PURI.*

Swingle, H.S., (1957). Biological means of increasing productivity in ponds. *FAO Fish Rep.***4** : 243 – 257.

Sinha, V.R.P., Gupta, M.V. Khan, D.P. chakraborty and M.V. Gupta. (1980). Preliminary observations on nitrogen balance of some ponds under composite fish culture. *Aquaculture Hungarica.* **2**:105 – 116.

Sinha, V.R.P. Gupta, M.V., Banerjee, M.K. and Kumar, D.(1973). Composite fish culture at kalyani, West Bengal. *J. Inland Fish Fish Soc. India.* **5**: 201 – 207.

Smith, J.H; Allison, F.E and Soulides, D.A., 1961. Evaluation of phosphobacteria as a soil inoculants. *Soil Sci. Soc. Am. Proc;* **25**; 109 – 111.

Shilo, M. and Rimon, A; 1982. Bamidgeh, **4**: 101-114

Schuetzle, D; Koskinen.and Horsfall 111, F.L., 1982. Chemometric modeling of waste water treatment process. J. Watert Pollute. Contron Fed;**54**: 457-465

Sun Yao and He Zhaoyang. 1997. Shrimp pond settlement- the quantity of nutrients dispersion and Seasonal changes between waterand mud contacting surfaces. Mar. Fish.Res. !8(!0 : 60-66.

Samocha, T.M; Lawrence, A.L; Horowite, A; Horowitz, S; 1998. The use of commercial probiotics in the production of of marine shrimpunder no water exchange. The Second International Conference on Recirculating Aquaculture. July 1998. Virginia polytechnic Institute, pp. 373-375.

Treer, T., Anicic, I., and Safmer, R. (1995). The growth and condition of common carps (*cyprinus carpio*) introduced into creation Vransko lake. *RIBARSTVO.* **53** (2): 63 – 73.

Tucker, Cs, Lloyd, S.W, 1985. Evaluation of a commercial bacterial amendment for improving water quality in channel catfish ponds, Mississiippi Agricultural and Forestry Experimental Station, Mississippi State University, Res. Rep. 10, 1-4.

Weaver, R.W.; and E,P. Dunigan, J.R. Parr, A.E. Hiltbold. 1974. Effect of two soil activators on crop yields activities of soil micro organisms in the southern United states. Southern Cooperative Serise Bulletin 189, Texas Aqricultural Experiment station, college Station, TX.

Walther, U. (1977). Fertilization of warm water fish ponds in Europe. FAP Fish Rep. **3** (44): 68 – 81.

Zhu,Yun, Yejin, Wan, Junhua, Hua, Dan, Mathias, H.A. (1994). The effect of manure Application rate and frequency upon fish yield in integrated fish farm ponds. The second paper of basic integrated fish farming theory and fishery bioeconomic research.

Zongheyangyo jichulilun he yuyeshengwu jianjiyanjiu lunwenji Beijing china. *Science press.*:128 – 159.

Zhu,Yun, Yang, Yejin, Wan, Junhua, Hua, Dan, Mathias, J.A. (1990). The effect of manure application rate and frequency upon fish yield in integrated fish farm ponds. *Aquacultures.* **91** (3-4) : 2333 – 251.

Zhang Zhongyang and Tang Yujun. 1989. Pond fish culture in China. Beijing Science publishing House. P. 45 – 47.

APHA 1995. Standard Methods for the Examination of Water and Wastewater . 19th Edition . American Public Health Association, Washington, DC, 1038p.

Austin,B. 1990.Methods in Aquatic, Bacteriology. John Wiley and Sons, New York, 425p.

Bauchop, T.& Elsden, S.R. 1960. The growth of microorganisms in relation to their energy supply . J. Gen. Micro.23:457-469.

Bhakta, J.N.; Bandyopadhyay , P.K.& Jana ,B.B. 2006 Effect of different doses of mixed fertillizer on some biogeochemical cycling bacterial population in carp culture pond. Tur. J. of Fish. and Aqu. Sc. 6:163-169.

Bhakta,J.N.2003. Fertilizer-microbial interactions in wastewater system: Influence of fertilizer dose and stocking density of fish. Ph . D. Thesis . University of Kalyani, Kalyani. 188p.

Bhakta,J.N.2006. Nutrient value of excreta of some freshwater fish species. Env. And Ecol.24S(2):370-374.

Bianchi, M.1998. Nouvelles approaches d'etude des reseaux microbiens. Annuals Limnol. 34:465-473. Billen, G. 1984. Heterotrophic utilization and regeneration nitrogen. In : Hobbie, J.E. & Williams, B.P.(eds.) Heteotrotrophic activity in the Sea. Plenum, New York.313-355 pp.

Boyd, C.E.1995. Bottom Solis, Sediment, and Pond Aquaculture. Chapman and Hall, New York. 348p.

Cho, B.C. & Azam, F. 1988. Major role of bacteria in biogeochemical fluxes in the ocean's interior. Nature 332:441-443.

Dawes. E.S.1985. Starvtion, survival and energy reserves. In: Fletcher, M.& Foodgate, G.D. (eds.) bacteria in Their Natural Environment. Academic, New York. 43-79 pp.

Diana, J.S & Fast, A.W. 1989. The effect of water exchange rate and density on yield of the walking catfish, *Clarias fuscus.* Aquaculture 78:267-276.

Ducklow, H.W. & Carlson, C.A.1992. Oceanic bacterial production. Adv. Microb. Ecol.12:113-181.

Gaur, A.C.;Dargan, K.S. & Neelakantan, K.S. 1995.Organic Manure. Publication and Information Division, Indian Conucil of Agricultural Research, New Delhi, Indian, 159p.

Giorgio, P.A.D. & Cole, J.J. 1998. Bactieral growth efficiency in natural aquatic systems. Annual Rev.of Ecol. And Syst. 29:503-541.

Gomez, K.A. & Gomez, A.A. 1984. Statistical Procedures for Agricultural Research 2nd Education. John Wiley and Sons, New York, 680p.

Jana, B.B.; Chakrabarti, P.; Biswas, J.K . &Ganguly, S.2001. Biogeochemical cycling bacteria as indices of pond fertilization: Importance of CNP ratios of input fertilizers. J.of Appl. Micro. 90:733-740.

Jana, B.B. & De, U.K.1990. Spatial and seasonal distribution of heterotrophic bacteria in pond water and sediments under different management practices. Int. Rev. der Ges. Hyd. 75:639-648.

Jana, B.B. & De, U.K. 1993. Management induced variability of the bacterioplanktom in fish farming ponds. J.of Aqu. In the Trop. 8:131-140.

Jana, B.B. & Patel, G.N. 1984. Spatial and seasonal variations of phosphate solubilizing bacteria in fish ponds of varying fish farming managements. Arch. Fer Hyd. 101:555-568.

Jana, B.B. & Patel, G.N. 1985. Distribution Pattern denitrifying bacteria in fish ponds of differing farming managements. Arch. Fur Hyd . 103:291-303.

Jana, B.B. & Patel,G.N. 1990. Pattern and magnitude of nitrate reduction in fish ponds differing in management practices. Aqua. Hugh. 6:53-63.

Jana, B.B. & Roy, S.K. 1985a. Distribution patterns of protein mineralizing and ammonifying bacterial populations in fish farming ponds under different management systems. Aquaculture 44:57-65.

Jana, B.B. & Roy, S.K.1985b. spatial and temporal changes of nitrifying bacterial population in fish ponds of different management practices. J.of Appl. Bact. 59:195-204.

Jana, B.B. & Roy, S.K.1986. Seasonal and spatial distribution Patten of nitrogen fixing bacteria in fish ponds under different management systems. Hydrobiology 137:45-54.

Milstein, A.; Azim, M.E,: Abdul, W.M. & Verdegem, M.C.J. 2003.The Effects of periphyton, fish and fertilizer dose on biological processes affecting water quality in earthen fish pond. Evn. Biol. Of Fish. 64:247-260.

Monod, J. 1942. Recherches sur la criossance des cultures bact eriennes. Hermann and
Co, Paris. Parson, T.; Baath, E.; Calrhom, M.; Lundkvisyt, H.;soderstrom, B.& Sohlenius, B. 1980. Trophic structure, biomass dynamics and carbon metabolism of soil organisms in Scots pine forest. Ecol. Bull. 32:419-462.

Pomeroy, L.R.; Wiebe, W.J.: Deibel, D.; Thompson, R.J. & Rowe, G.T. 1991. Bacterial responses to temperature and substrate concentration during the Newfoundland spring bloom Mar. Ecol. Prog. Ser. 75:143-159.

Rodina, A.G.1972. Methods in Aquatic Microbiology. In: Colwell, R.R. & Zambruski, M.S. (eds) University Park Press, Baltimore, Butterworths, London, 461p. \

Vollenweider, R.A. 1974. A Manual on Methods for Measuring Primary Production in Aquatic Environments. Blackwell Scientific Publications, Oxford, 225p.

Williams, P.B. 1981. Microbial contribution to overall marine plankton metabolism: direct measurements of respiration. Oceanol. Acta 4:359-364